Redden Island Bird Checklist

Australian Brush turkov	D	Nankeen Kestrel	R
Australian Brush-turkey			
Orange-footed Scrubfowl		Brown Falcon	R
Brown Quail		Peregrine Falcon	R
Magpie Goose		Bush Stone-curlew	U
Radjah Shelduck		Beach Stone-curlew	R
Green Pygmy Goose		Australian Pied Oystercatcher	U
Grey Teal		Black-winged Stilt	U
Pacific Black Duck		Pacific Golden Plover	U,Mn
Australasian Grebe		Red-capped Plover	С
Spotted Dove	_	Double-banded Plover	R,Mnz
Peaceful Dove	C	Lesser Sand Plover	C,Mn
Bar-shouldered Dove	_	Greater Sand Plover	C,Mn
Superb Fruit-Dove		Masked Lapwing	С
Rose-crowned Fruit-Dove	С	Black-tailed Godwit	R,Mn
Pied Imperial-Pigeon	C,Mn	Bar-tailed Godwit	C,Mn
Topknot Pigeon		Whimbrel	С
Papuan Frogmouth		Eastern Curlew	С
Large-tailed Nightjar	U,h	Terek Sandpiper	C,Mn
Australian Swiftlet	С	Common Sandpiper	U,Mn
White-throated Needletail	R,Mn	Grey-tailed Tattler	U,Mn
Fork-tailed Swift	R,Mn	Wandering Tattler	Ć,Mn
Lesser Frigatebird		Marsh Sandpiper	Ć,Mn
Greater Frigatebird	R	Common Greenshank	Ć,Mn
Australasian Darter	U	Ruddy Turnstone	U,Mn
Little Pied Cormorant		Great Knot	C,Mn
Little Black Cormorant	R	Red-necked Stint	C,Mn
Pied Cormorant	R	Sharp-tailed Sandpiper	C,Mn
Australian Pelican	C	Curlew Sandpiper	U,Mn
Black-necked Stork	Ū	Ruff	V
Eastern Great Egret	Ċ	Little Tern	C,Mn
Intermediate Egret		Gull-billed Tern	C
Great-billed Heron	_	Caspian Tern	C
Striated Heron		Common Tern	U
White-faced Heron	_	Lesser Crested Tern	
Little Egret	_	Crested Tern	R C
Eastern Reef Egret		Silver Gull	C
Nankeen Night-Heron	D	Red-tailed Black-Cockatoo	C
Australian White Ibis	C	Sulphur-crested Cockatoo	C C
Straw-necked Ibis	U	Rainbow Lorikeet	Ċ
Royal Spoonbill	R	Scaly-breasted Lorikeet	Ċ
Eastern Osprey	C	Little Lorikeet	R
Black-shouldered Kite	R	Double-eyed Fig-Parrot	C
Pacific Baza		Red-winged Parrot	R
	R	Pheasant Coucal	C
White-bellied Sea-Eagle	U	Eastern Koel	U,Mn,h
Whistling Kite	U	Channel-billed Cuckoo	U,Mn
Brahminy Kite	С	Horsfield's Bronze-Cuckoo	R
Black Kite	U	Little Bronze-Cuckoo	Ċ
Brown Goshawk	Ū	Pallid Cuckoo	R
Grey Goshawk	R	Fan-tailed Cuckoo	R
		. a tanea cacitoo	11

Brush Cuckoo	C,h	Rufous Whistler	U
Oriental Cuckoo	R,Mn	Little Shrike-thrush	R
Azure Kingfisher	R	Australasian Figbird	Ċ
Little Kingfisher	R	Yellow Oriole	C
Buff-breasted Paradise-Kingfisher	R,Mn	Olive-backed Oriole	R
Laughing Kookaburra	С	White-breasted Woodswallow	C
Forest Kingfisher	С	Black Butcherbird	Č
Sacred Kingfisher	С	Spangled Drongo	C
Collared Kingfisher	С	Rufous Fantail	U
Rainbow Bee-eater	С	Grey Fantail	Ü
Dollarbird	C,Mn	Northern Fantail	Ü
Noisy Pitta	U	Willie Wagtail	C
Great Bowerbird	R	Torresian Crow	U
Red-backed Fairy-wren	U	Leaden Flycatcher	C
Lovely Fairy-wren	U	Satin Flycatcher	R
Large-billed Scrubwren	R	Shining Flycatcher	C
Large-billed Gerygone	С	White-eared Monarch	R
Fairy Gerygone	С	Black-faced Monarch	Ü
Striated Pardalote	U	Spectacled Monarch	C
Yellow-spotted Honeyeater	С	Magpie-lark	C
Graceful Honeyeater	С	Mangrove Robin	U
Varied Honeyeater	U	Tawny Grassbird	R
Yellow Honeyeater	С	Silvereye	U
Brown-backed Honeyeater	С	Welcome Swallow	C
Dusky Honeyeater	С	Fairy Martin	C
Brown Honeyeater	C	Tree Martin	U
White-throated Honeyeater	C	Metallic Starling	C
Helmeted Friarbird	С	Common Myna	C
Noisy Friarbird	R C	Mistletoebird	C
Black-faced Cuckoo-shrike White-bellied Cuckoo-shrike	C	Olive-backed Sunbird	C
Barred Cuckoo-shrike	Ü	Double-barred Finch	_
Cicadabird	C,h	Crimson Finch	R
White-winged Triller	R	Nutmeg Mannikin	R
Varied Triller	C	Chestnut-breasted Mannikin	С
Grey Whistler	IJ	House Sparrow	U
C. C, William			U
		•	

LIST ANNOTATIONS

C=Common Likely to be recorded on most visits
U=Uncommon Recorded regularly but not every visit
R=Rare Recorded infrequently
V=Vagrant Does not usually occur in Australia

h=More often heard than seen
Mn=Migrant from the North (approx Oct-Mar)
Mw =Migrant from the West (approx May-Sep)

Vegetation Types

- Tall woodland with Variable-barked Bloodwood *Corymbia erythrophloia* & Moreton Bay Ash *C. tessellaris* (15m)
- Tall open woodland with Pink Bloodwood *C. intermedia* (30m)
- Tall woodland 15-18m with Broad-leaved Tea Tree *Melaleuca leucadendra*
- A small area of Tall closed forest with Beach Tamarind Cupaniopsis anacardioides, White Tamarind C. foveolata, Beach Cherry Eugenia reinwardtiana and Supplejack Flagellaria indica
- Cannonball Mangrove Xylocarpus granatum + Slender-fruited Mangrove Bruguiera parviflora
- Rhizophora spp + Grey Mangrove Avicenia marina
- A high density *Ceriops spp.* stand
- A beach community
- A highly mixed fringing community

Habitat data courtesy of Cairns Port Authority 'Environmental Baseline Studies Volume Two'

Definition: The C/U/R annotations in this list give an indication of how easy or difficult it might be to find a species. They indicate the likelihood that an experienced birder might record the species, during a day bird-watching, at the right time of year, in the appropriate habitat.

Updated April 2013

© BirdLife Northern Queensland

WHO ARE WE?

BirdLife Northern Queensland

is a branch of the national organisation BirdLife Australia, formed in 2012 by the merger of Birds Australia and BOCA.

BirdLife Northern Queensland organizes both scientific and social activities. These include bird surveys, outings, workshops and presentations with guest speakers. The regular field trips and campouts cater for all levels of expertise from beginner to professional and emphasise fun as well as worthwhile conservation projects.

Members receive a quarterly newsletter *Contact Call* detailing local activities and also enjoy *BirdLife Australia* magazine dealing with national bird issues.

You can join BirdLife Australia and help the conservation of Australian birds at www.birdlife.org.au

New or unusual sightings?

Register with Eremaea Birds

www.eremaea.com

INFORMATION

Redden Island is situated in the Cairns suburb of Machans Beach, just north of Cairns International Airport. About 68ha of Redden Island is owned by the Cairns Port Authority. Surrounded by the Barron River, Redden Creek and the Coral Sea, it contains a high diversity of habitats in a relatively small area and attracts a large number of bird species.


Waders (shorebirds) are mainly found at the mouth of the Barron River on the sandbars and are best viewed on an incoming tide. At higher tide they move to the opposite bank of the river, on Casuarina Point. Other bird species may be found by following the various walking tracks around the island. The Great-billed Heron and Little Kingfisher might be sighted on Redden Creek and the Mangrove Robin is hard to find along the Barron River opposite the airport.

Migrants (Sept-April) include Pied Imperial Pigeon, Oriental Cuckoo, Dollarbird and most waders. The Red-winged Parrot is unusual on the coast and the Great Bowerbird was the first sighting in Cairns.

Access and Facilities

Vehicle access is via the Marshall St bridge over Redden Creek to Redden Island, and Cinderella St which leads to the Barron River mouth. Walking tracks traverse most habitats and are accessed via 3 pedestrian gates along Cinderella St.

Bring suitable walking shoes and insect repellent



REDDEN ISLAND

Bird Checklist No 2



BirdLife Northern Queensland



Email northernqld@birdlife.org.au
Web www.birdsaustralianq.org
www.birdlife.org.au
Facebook Birdlife Northern Queensland